

# A Form of the Borel–Cantelli Lemma

NADJIB BOUZAR

*Department of Mathematics, Northwestern University,  
Evanston, Illinois 60201*

## INTRODUCTION

In this paper we prove a strong version of the Borel–Cantelli lemma. Our proof is based on a convergence theorem in martingale theory and a local convergence theorem for sequences of positive random variables [5]. We then apply our result (Theorem 2) to deduce both a form of the Borel–Cantelli lemma due to Dubins and Freedman [1], and a more recent result due to Hill [2]. We also prove a strong law of large numbers for sequences of positive random variables (Theorem 4), and as an application we prove a convergence theorem on infinite series that is a generalization of a theorem of Dini [2, 3].

## 1. PRELIMINARIES

Let  $(\Omega, \mathcal{F}, P)$  be a probability space and  $(\mathcal{F}_n, n \geq 1)$  an increasing sequence of sub- $\sigma$ -fields of  $\mathcal{F}$ . A sequence  $(X_n, n \geq 1)$  of random variables is said to be adapted if for each  $n \geq 1$ ,  $X_n$  is  $\mathcal{F}_n$ -measurable. For  $p \in [1, +\infty)$  we denote by  $L^p(\Omega, \mathcal{F}, P)$  the space of random variables of  $p$ th integrable power. For an adapted sequence of integrable random variables  $(X_n, n \geq 1)$  we define  $(S_n, n \geq 1)$  by,

$$S_n = \sum_{i=1}^n X_i.$$

We define  $(s_n, n \geq 1)$  by,

$$\begin{aligned} s_1 &= 0 \\ s_{n+1} &= \sum_{i=1}^n E(X_{i+1} | \mathcal{F}_i), \quad n \geq 1. \end{aligned}$$

If moreover  $(X_n, n \geq 1)$  is in  $L^p(\Omega, \mathcal{F}, P)$  for some  $p \in [1, +\infty)$  we define  $(A_n^{(p)}, n \geq 1)$  by

$$A_1^{(p)} = 0$$

$$A_{n+1}^{(p)} = \sum_{i=1}^n E(|X_{i+1}|^p | \mathcal{F}_i), \quad n \geq 1.$$

## 2. CONVERGENCE RESULTS

We start out with a lemma.

LEMMA 1. *Let  $(X_n, n \geq 1)$  be an adapted sequence of positive random variables in  $L^p(\Omega, \mathcal{F}, P)$  for some  $p \in [1, +\infty)$ . Let  $f: [0, \infty) \rightarrow \mathbb{R}$  be a strictly positive and increasing function such that  $\int_0^{+\infty} f^{-p}(x) dx < \infty$ . Then,*

$$\sum_{i=1}^{+\infty} f^{-p}(A_{i+1}^{(p)}) E(|X_{i+1} - E(X_{i+1} | \mathcal{F}_i)|^p | \mathcal{F}_i) < \infty \quad \text{a.s.}$$

*Proof.* For  $i \geq 1$ ,  $|X_{i+1} - E(X_{i+1} | \mathcal{F}_i)|^p \leq 2^p(|X_{i+1}|^p + |E(X_{i+1} | \mathcal{F}_i)|^p)$ . Therefore by Jensen's inequality, by the definition of  $A_i^{(p)}$  and by the fact that  $X_i \geq 0$  for each  $i \geq 1$ , we have

$$E(|X_{i+1} - E(X_{i+1} | \mathcal{F}_i)|^p | \mathcal{F}_i) \leq 2^{p+1} E(X_{i+1}^p | \mathcal{F}_i)$$

$$= 2^{p+1} (A_{i+1}^{(p)} - A_i^{(p)}).$$

Hence it is enough to show that

$$\sum_{i=1}^{+\infty} f^{-p}(A_{i+1}^{(p)}) (A_{i+1}^{(p)} - A_i^{(p)}) < \infty \quad \text{a.s.}$$

But the fact that  $f$  is increasing implies that

$$\sum_{i=1}^{+\infty} f^{-p}(A_{i+1}^{(p)}) (A_{i+1}^{(p)} - A_i^{(p)}) \leq \sum_{i=1}^{+\infty} \int_{A_i^{(p)}}^{A_{i+1}^{(p)}} f^{-p}(x) dx$$

$$\leq \int_0^{+\infty} f^{-p}(x) dx < \infty.$$

We now prove the main theorem.

THEOREM 2. *Let  $(X_n, n \geq 1)$  be an adapted sequence of random variables such that  $0 \leq X_n \leq 1$  for each  $n \geq 1$ . Let  $f: [0, \infty) \rightarrow \mathbb{R}$  be a strictly*

positive and increasing function such that  $\int_0^{+\infty} f^{-p}(x) dx < \infty$  for some  $p \in [1, 2]$ . Then

$$\lim_{n \rightarrow +\infty} \frac{S_n - s_n}{f(s_n)} < \infty \quad \text{a.s.}$$

The limit is equal to 0 on the set  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ .

*Proof.* We first prove that  $\lim_{n \rightarrow +\infty} [(S_n - s_n)/f(s_n)] = 0$  a.s. on the set  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ . Let  $M_n = S_n - s_n$  for each  $n \geq 1$ . Since  $M_{n+1} - M_n = X_{n+1} - E(X_{n+1} | \mathcal{F}_n)$  ( $M_n, n \geq 1$ ) is a martingale. Moreover, since  $p \geq 1$  and  $0 \leq X_i \leq 1$  for each  $i \geq 1$ ,

$$A_{i+1}^{(p)} = \sum_{j=1}^i E(X_{j+1}^p | \mathcal{F}_j) \leq \sum_{j=1}^i E(X_{j+1} | \mathcal{F}_j) = s_{i+1}.$$

Hence,  $f^{-p}(s_{i+1}) \leq f^{-p}(A_{i+1}^{(p)})$  and by Lemma 1,

$$\sum_{i=1}^{+\infty} f^{-p}(s_{i+1}) E(|M_{i+1} - M_i|^p | \mathcal{F}_i) < \infty \quad \text{a.s.}$$

Now let  $W_1 = 0$ , and for  $n \geq 1$   $W_{n+1} = \sum_{i=1}^n f^{-1}(s_{i+1})(M_{i+1} - M_i)$ . For every  $n \geq 1$ ,  $f(s_{n+1})$  is  $\mathcal{F}_n$ -measurable. Hence  $(W_n, n \geq 1)$  is a martingale and  $\sum_{i=1}^{+\infty} E(|W_{i+1} - W_i|^p | \mathcal{F}_i) < \infty$  a.s. Therefore by Corollary 2.8.5 [4, p. 67],  $W_n$  converges a.s. Since  $\lim_{x \rightarrow +\infty} f(x) = +\infty$  and  $f$  is increasing,  $[\lim_{n \rightarrow +\infty} f(s_n) = +\infty] = [\lim_{n \rightarrow +\infty} s_n = +\infty]$ . Hence by Kronecker's lemma  $\lim_{n \rightarrow +\infty} [M_n/f(s_n)] = 0$  a.s. on  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ . To prove the convergence on the set  $[\lim_{n \rightarrow +\infty} s_n < \infty]$  we apply Theorem 2 in [5]. It says that on the set  $[\sup_{n \geq 1} s_n < \infty]$ ,  $S_n$  converges a.s. if and only if  $s_n$  converges a.s. Therefore  $S_n$  converges a.s. on  $[\lim_{n \rightarrow +\infty} s_n < \infty]$ . Moreover the fact that  $f$  is increasing implies that  $f(s_n)$  converges on  $[\lim_{n \rightarrow +\infty} s_n < +\infty]$ . Hence  $(S_n - s_n)/f(s_n)$  converges on  $[\lim_{n \rightarrow +\infty} s_n < \infty]$ .  
Q.E.D.

Next we use Theorem 2 to deduce the versions of the Borel-Cantelli lemma given in [1, 2]. The sequence  $(X_n, n \geq 1)$  will be as in Theorem 2, and for  $k \geq 1$  we let  $\log_k x$  be the  $k$ th iterated logarithm of  $x$  (e.g.,  $\log_3 x = \log \log \log x$ ).

COROLLARY 3. (i) For every integer  $k \geq 1$  and for every  $p \in (1, 2]$ ,

$$\lim_{n \rightarrow +\infty} \frac{S_n - s_n}{(s_n \log_1(s_n) \cdots \log_{k-1}(s_n) \log_k^p(s_n))^{1/p}} = 0$$

a.s. on  $\left[ \lim_{n \rightarrow +\infty} s_n = +\infty \right]$ .

(ii) If  $f: [0, \infty) \rightarrow \mathbb{R}$  is a strictly positive and increasing function such that  $\lim_{x \rightarrow +\infty} [x/f(x)] = a < \infty$  then  $S_n/f(s_n)$  converges a.s. to a finite limit. The limit is equal to  $a$  on the set  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ .

*Proof.* (i) For  $k \geq 1$  we define  $f_k: [a_k, +\infty) \rightarrow \mathbb{R}$  by  $f_k(x) = (x \log_1 x \cdots \log_{k-1} x \log_k^p x)^{1/p}$  where  $a_k$  is taken to be large enough so that  $f_k$  is well defined, strictly positive and increasing. Let  $f_k^*: [0, \infty) \rightarrow \mathbb{R}$  be the extension of  $f$  that is identically equal to  $f_k(a_k)$  on the interval  $[0, a_k]$ ;  $f_k^*$  is strictly positive and increasing. Moreover,

$$\int_{a_k}^{+\infty} f_k^{-p}(x) dx = ((p-1) \log_k^{p-1}(a_k))^{-1} < \infty.$$

Therefore  $\int_0^{+\infty} (f_k^*(x))^{-p} dx < \infty$ . Hence, by Theorem 2,

$$\lim_{n \rightarrow +\infty} \frac{S_n - s_n}{f_k^*(s_n)} = \lim_{n \rightarrow +\infty} \frac{S_n - s_n}{f_k(s_n)} = 0$$

a.s. on  $\left[ \lim_{n \rightarrow +\infty} s_n = +\infty \right]$ .

(ii) The fact that  $\lim_{x \rightarrow +\infty} [x/f(x)]$  exists and is finite implies that  $\int_0^{+\infty} f^{-2}(x) dx < \infty$ . Therefore, by Theorem 2,  $(S_n - s_n)/f(s_n)$  converges a.s. We have a.s. on the set  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ ,

$$\begin{aligned} \lim_{n \rightarrow +\infty} \frac{S_n}{f(s_n)} &= \lim_{n \rightarrow +\infty} \left( \frac{S_n - s_n}{f(s_n)} + \frac{s_n}{f(s_n)} \right) \\ &= \lim_{x \rightarrow +\infty} \frac{x}{f(x)} = a. \end{aligned}$$

On the set  $[\lim_{n \rightarrow +\infty} s_n < \infty]$ ,  $f(s_n)$  converges because  $f$  is increasing, therefore  $S_n/f(s_n)$  converges a.s. Q.E.D.

*Remarks.* (1) Part (ii) of Corollary 3 was proved by Dubins and Freedman [1] for  $f(x) = x + EX_1$  with  $EX_1 > 0$ . Part (i) is due to Hill [2].

(2) It is an easy exercise to generalize Theorem 2 to an adapted sequence of nonnegative uniformly bounded random variables.

(3) Suppose that  $f$  satisfies the following condition: for any two sequences of real numbers  $(a_n, n \geq 1)$  and  $(b_n, n \geq 1)$  such that  $\lim_{n \rightarrow +\infty} (a_n/b_n) = 1$ ,  $\lim_{n \rightarrow +\infty} [f(a_n)/f(b_n)] < \infty$ . Then under the assumptions of Theorem 2 (and  $EX_1 > 0$ ),  $(S_n - s_n)/f(s_n)$  converges a.s. to a finite limit. The limit is equal to 0 on  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ . The convergence on  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$  follows from Theorem 2 and Remark 1 above. On  $[\lim_{n \rightarrow +\infty} s_n < \infty]$  we use the same argument as in the proof of Theorem 2.

The functions  $(f_k, k \geq 1)$  in the proof of Corollary 3 satisfy the condition above.

Next we prove a strong law of large numbers for sequences of positive random variables. As an application we prove a result on infinite series that is a simultaneous generalization of a theorem of Dini [3, p. 290], and Lemma 1 in [2].

**THEOREM 4.** *Let  $(X_n, n \geq 1)$  be an adapted sequence of positive integrable random variables. Let  $f: [0, +\infty) \rightarrow \mathbb{R}$  be a strictly positive and increasing function such that  $\int_0^{+\infty} f^{-1}(x) dx < \infty$ . Then,*

- (i)  $\sum_{n=1}^{+\infty} [X_n/f(s_n)] < \infty$  a.s.
- (ii)  $\lim_{n \rightarrow +\infty} [S_n/f(s_n)] = 0$  a.s. on  $[\lim_{n \rightarrow +\infty} s_n = +\infty]$ .

*Proof.* (i) For each  $n \geq 1$ , let  $W_n = \sum_{i=1}^n f^{-1}(s_i) X_i$ ;  $(W_n, n \geq 1)$  is adapted and positive. Moreover, since  $f(s_{n+1})$  is  $\mathcal{F}_n$ -measurable and  $f$  is increasing,

$$\begin{aligned} 0 \leq E(W_{n+1} - W_n | \mathcal{F}_n) &= E(f^{-1}(s_{n+1}) X_{n+1} | \mathcal{F}_n) \\ &= f^{-1}(s_{n+1})(s_{n+1} - s_n) \\ &\leq \int_{s_n}^{s_{n+1}} f^{-1}(x) dx. \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{n=1}^{+\infty} E(W_{n+1} - W_n | \mathcal{F}_n) &\leq \sum_{n=1}^{+\infty} \int_{s_n}^{s_{n+1}} f^{-1}(x) dx \\ &\leq \int_0^{\infty} f^{-1}(x) dx. \end{aligned}$$

Therefore

$$\sum_{n=1}^{+\infty} E(W_{n+1} - W_n | \mathcal{F}_n) < \infty \quad \text{a.s.}$$

Applying Theorem 2 in [5] we conclude that  $W_n$  converges a.s.

- (ii) Since  $\lim_{x \rightarrow +\infty} f(x) = +\infty$  and  $f$  is increasing,

$$\left[ \lim_{n \rightarrow +\infty} f(s_n) = +\infty \right] = \left[ \lim_{n \rightarrow +\infty} s_n = +\infty \right].$$

The conclusion follows by using (i) and Kronecker's lemma.

Q.E.D.

COROLLARY 5. Let  $(d_n, n \geq 1)$  be a sequence of positive numbers with  $D_n = \sum_{i=1}^n d_i$ . Let  $f: [0, \infty) \rightarrow \mathbb{R}$  be a strictly positive and increasing function such that  $\int_0^{+\infty} f^{-1}(x) dx < \infty$ . Then

$$\sum_{n=1}^{+\infty} \frac{d_n}{f(D_n)} < \infty.$$

*Proof.* Without loss of generality we may assume  $d_1 = 0$ . Let  $X_n = d_n$  for each  $n \geq 1$ . Therefore  $S_n = s_n = D_n$ . The conclusion follows by applying Theorem 4(i) to the sequence  $(X_n, n \geq 1)$ . Q.E.D.

### REFERENCES

1. L. DUBINS AND D. FREEDMAN, A sharper form of the Borel–Cantelli lemma and the strong law, *Ann. Math. Stat.* **36** (1965), 800–807.
2. T. P. HILL, A stronger form of the Borel–Cantelli lemma, *Illinois J. Math.* **27** (1983), 240–243.
3. K. KNOPP, “Theory and Applications of Infinite Sums,” Blackie, Glasgow, 1946.
4. W. F. STOUT, “Almost Sure Convergence,” Academic Press, New York, 1974.
5. D. Q. TUYËN, On the asymptotic behavior of random variables. *Ann. Inst. H. Poincaré* **XVII** (1981), 63–73.